


FORMULE D'ADDITIONE E SOTTRAZIONE



Fixarsi $P(\cos \alpha; \sin \alpha)$
 $Q(\cos \beta; \sin \beta)$.

$$R(\cos(\alpha - \beta); \sin(\alpha - \beta))$$

$$A(1, 0)$$

$$\triangle OAR \cong \triangle QOP$$

Per costruzione $\overline{QP} = \overline{AR}$ quindi:

$$\sqrt{(\cos \alpha \cdot \cos \beta)^2 + (\sin \alpha \cdot \sin \beta)^2} = \sqrt{(\cos(\alpha - \beta) - 1)^2 + \sin^2(\alpha - \beta)}$$

$$\boxed{\cos^2 \alpha + \cos^2 \beta - 2 \cos \alpha \cos \beta \cdot \sin^2 \alpha + \sin^2 \beta - 2 \sin \alpha \sin \beta} = \cos^2(\alpha - \beta) + 1 - 2 \cos(\alpha - \beta) + \sin^2(\alpha - \beta)$$

$$1 + 1 - 2(\cos \alpha \cos \beta + \sin \alpha \sin \beta) = 1 + 1 - 2 \cos(\alpha - \beta)$$

$$\cancel{2} - 2(\cos \alpha \cos \beta + \sin \alpha \sin \beta) = \cancel{-2} \cos(\alpha - \beta)$$

$$\boxed{\cos(\alpha - \beta) = \cos \alpha \cos \beta + \sin \alpha \sin \beta} \quad (*) \quad (1)$$

||

$$\cos(\alpha + \beta) = \cos(\alpha - (-\beta)) = \cos \alpha \cos(-\beta) + \sin \alpha \sin(-\beta) =$$


$$= \cos \alpha \cos \beta + \sin \alpha (-\sin \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta.$$

$$\boxed{\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta} \quad (*) \quad (2)$$

$$\sin(\alpha + \beta) = \cos \left[\frac{\pi}{2} - (\alpha + \beta) \right] = \cos \left[\frac{\pi}{2} - \alpha - \beta \right]$$

$$= \cos \left[\left(\frac{\pi}{2} - \alpha \right) - \beta \right] = \cos \left(\frac{\pi}{2} - \alpha \right) \cos \beta + \sin \left(\frac{\pi}{2} - \alpha \right) \sin \beta =$$

$$= \sin \alpha \cos \beta + \cos \alpha \sin \beta$$



$$\boxed{\sin(\alpha + \beta) = \sin \alpha \cos \beta + \cos \alpha \sin \beta} \quad (*) \quad (3)$$

$$\sin(\alpha - \beta) = \sin(\alpha + (-\beta)) = \sin \alpha \cos(-\beta) + \cos \alpha \sin(-\beta) =$$

$$= \sin \alpha \cos \beta - \cos \alpha \sin \beta$$

$$\boxed{\sin(\alpha - \beta) = \sin \alpha \cos \beta - \cos \alpha \sin \beta} \quad (*) \quad (4)$$

$$\operatorname{tg}(\alpha + \beta) = \frac{\operatorname{sen}(\alpha + \beta)}{\cos(\alpha + \beta)} = \frac{\operatorname{sen}\alpha \cos\beta + \cos\alpha \operatorname{sen}\beta}{\cos\alpha \cos\beta - \operatorname{sen}\alpha \operatorname{sen}\beta} = (\cdot)$$

dividiamo numeratore e denominatore per $\cos\alpha \cos\beta \neq 0$;
 allora supponiamo $\cos\alpha \neq 0 \Rightarrow \alpha \neq \frac{\pi}{2} + k\pi \quad k \in \mathbb{Z}$ e $\cos\beta \neq 0 \Rightarrow$

$$\boxed{\beta \neq \frac{\pi}{2} + h\pi \quad \text{casi } h \in \mathbb{Z}}$$

$$(\cdot) = \frac{\frac{\operatorname{sen}\alpha \cos\beta}{\cos\alpha \cos\beta} + \frac{\cos\alpha \operatorname{sen}\beta}{\cos\alpha \cos\beta}}{\frac{\cos\alpha \cos\beta}{\cos\alpha \cos\beta} - \frac{\operatorname{sen}\alpha \operatorname{sen}\beta}{\cos\alpha \cos\beta}} = \frac{\operatorname{tg}\alpha + \operatorname{tg}\beta}{1 - \operatorname{tg}\alpha \operatorname{tg}\beta}$$

$$\boxed{\operatorname{tg}(\alpha + \beta) = \frac{\operatorname{tg}\alpha + \operatorname{tg}\beta}{1 - \operatorname{tg}\alpha \operatorname{tg}\beta}} \quad (*) \quad (5)$$

$$\operatorname{tg}(\alpha - \beta) = \operatorname{tg}(\alpha + (-\beta)) = \frac{\operatorname{tg}\alpha + \operatorname{tg}(-\beta)}{1 - \operatorname{tg}\alpha \operatorname{tg}(-\beta)} = \frac{\operatorname{tg}\alpha - \operatorname{tg}\beta}{1 + \operatorname{tg}\alpha \operatorname{tg}\beta}$$

$$\boxed{\operatorname{tg}(\alpha - \beta) = \frac{\operatorname{tg}\alpha - \operatorname{tg}\beta}{1 + \operatorname{tg}\alpha \operatorname{tg}\beta}} \quad (*) \quad (6)$$

$$\operatorname{tg}(\alpha + \beta) = \frac{\cos(\alpha + \beta)}{\sin(\alpha + \beta)} = \frac{\cos \alpha \cos \beta - \sin \alpha \sin \beta}{\sin \alpha \cos \beta + \cos \alpha \sin \beta} = (\ldots)$$

Si dividono numeratore e denominatore per $\sin \alpha \sin \beta \neq 0$, allora
 $\sin \alpha \neq 0 \Rightarrow \alpha \neq k\pi, k \in \mathbb{Z}$ e $\sin \beta \neq 0 \Rightarrow \beta \neq h\pi$ con $h \in \mathbb{Z}$.

$$(\ldots) = \frac{\frac{\cos \alpha \cos \beta}{\sin \alpha \sin \beta} - \frac{\sin \alpha \sin \beta}{\sin \alpha \sin \beta}}{\frac{\sin \alpha \cos \beta}{\sin \alpha \sin \beta} + \frac{\cos \alpha \sin \beta}{\sin \alpha \sin \beta}} = \frac{\operatorname{tg} \alpha \operatorname{tg} \beta - 1}{\operatorname{tg} \beta + \operatorname{tg} \alpha}$$

$$\boxed{\operatorname{tg}(\alpha + \beta) = \frac{\operatorname{tg} \alpha \operatorname{tg} \beta - 1}{\operatorname{tg} \beta + \operatorname{tg} \alpha}}$$

*

(+)

$$\operatorname{tg}(\alpha - \beta) = \operatorname{tg}(\alpha + (-\beta)) = \frac{\operatorname{tg} \alpha \operatorname{tg}(-\beta) - 1}{\operatorname{tg}(-\beta) + \operatorname{tg} \alpha} =$$

$$= \frac{\operatorname{tg} \alpha (-\operatorname{tg} \beta) - 1}{-\operatorname{tg} \beta + \operatorname{tg} \alpha} = \frac{1 + \operatorname{tg} \alpha \operatorname{tg} \beta}{\operatorname{tg} \beta - \operatorname{tg} \alpha}$$

$\alpha \neq k\pi, k \in \mathbb{Z}$
 $\beta \neq h\pi, h \in \mathbb{Z}$

$$\boxed{\operatorname{tg}(\alpha - \beta) = \frac{1 + \operatorname{tg} \alpha \operatorname{tg} \beta}{\operatorname{tg} \beta - \operatorname{tg} \alpha}}$$

*

(8)

$$\sec(\alpha+\beta) = \frac{1}{\cos(\alpha+\beta)} = \frac{1}{\cos\alpha\cos\beta - \sin\alpha\sin\beta} =$$

$$= \frac{1}{\cos\alpha\cos\beta(1 - \tan\alpha\tan\beta)} = \frac{\sec\alpha\sec\beta}{1 - \sec\alpha\sec\beta \tan\alpha\tan\beta}$$

$$\tan\alpha = \sqrt{1 - \cos^2\alpha} = \sqrt{1 - \frac{1}{\sec^2\alpha}} = \frac{\sqrt{\sec^2\alpha - 1}}{\sec\alpha}$$

$$\cosec(\alpha+\beta) = \frac{1}{\sin(\alpha+\beta)} = \frac{1}{\sin\alpha\cos\beta + \cos\alpha\sin\beta} =$$

$$= \frac{1}{\sin\alpha\sin\beta\left(\frac{\cos\beta}{\sin\beta} + \frac{\cos\alpha}{\sin\alpha}\right)} = \frac{\cosec\alpha\cosec\beta}{\cot\beta + \cot\alpha}$$

$$\cot\beta = \cosec\beta \sqrt{1 - \sin^2\beta} = \cosec\beta \sqrt{1 - \frac{1}{\cosec^2\beta}}$$